Field Trip No. 27.

Trangie plots, and collecting Narromine – Canberra.

26-27/10/1951

K. H. L. Key

Field Trip No. 27 — 26-27/10/51

Transgic plots. and collecting Narromine – Canberra.

Farm plot. – Possible controls lie on the road side of the plot, where the low-lying area continues. One is situated a few hundred yards from the plot Nyale Nundle, large. – A good control lies immediately adjacent to the plot, on the west side. Others should exist along the line of contact between the self-mulching plain & light country to the south which extends over to the small plot and the large fenced area near the gate.

26/10/51

36749. Narromine. 760. Rocky hill with Callitris glauca & E. blakelyi, occ. phyllode wattle. Aristida & Stipa tussocks. Ced. (m) / 774. Dubbo. 780. Thinned from albens + (mainly) C. glauca on gentle rise. Pasture mainly Stipa tussocks to 1ft. Ced (m) Chat (m) Eckphantus (m) 792. Geurie. 799. Marysvale. 802. Gradly thinned from albens, melliodora & Angophora ? intermedia. Much Paterson's curse / Basalt. Brachyscama (m) 806. Mullington. Macrolophotia sp., froggattina 27.10.51 807. Left Mullington. 812. Albens var. woodland on australis, Cryptobothrus clinopthorus chocolate soil derived from basalt & limestone. Pasture to 9 in., largely weeds, a little bare space. froggattia quite numerous. Gonacea sp. 826. Thinned albens on Cryptobothrus chrysophorus, Apurga sp. slope with Stipa & Aristida tussocks leading down into depression with continuous low cover and tussock. Frogg., Brachyscama (m). 841. Narromine Lee.

3842. Stuartiana on steep slope with Poa tussock leading to flat with closed pasture to bin. with Tipuna sp. melliodora. Both trees considerably thinned. 848. Along Paratettix australis, T. argillaceus, Macrolophotia sp. 854. Rocky hillside with C. calcarata, & E. deabbata & Austrabris guttulosa, Pycnostictus sp. n., Aylota sp. blakelyi descending to depression with continuous pasture. Paratettix in depression. 862. Mauldra. Isonica sp. Cryptobothrus chrysophorus, Tipuna sp. 871. Cudal. 874. Thinned & cleared albens with Stipa pasture Pycnostictus sp. n. some bare ground. Brachyscama (?). lower ground with C. glauca & occ. woollsiana had no acrids. 874. Canowindra. 912. Cowra cemetery. Portion has been undisturbed for a long time, except for burning. Themeda evans believingly predominant, along with a stalky grass, Lomandra & the blue-flowered liliaceous tussock plant.